



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

100-1610

Date of Issuance:

1/3/17

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

FORCE EVO

Name and Address of Registrant (include ZIP Code):

Charles T. Levey
 Team Lead Insecticides/Fungicides
 Syngenta Crop Protection, LLC
 P.O. Box 18300
 Greensboro, NC 27419

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kable Bo Davis, Product Manager 03 and 04
 Invertebrate and Vertebrate Branch 1, Registration Division (7505P)

Date:

1/3/17

2. You are required to comply with the data requirements described in the DCI Order identified below:

- a. Tefluthrin GDCI-128912-1293

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI Order listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 100-1610.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 03/03/2016
- Alternate CSF 1 dated 05/26/2016

If you have any questions, you may contact Elizabeth Fertich at 703-347-8560 or via email at fertich.elizabeth@epa.gov.

Enclosure

**RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS**
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR
DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S
CERTIFICATION.

GROUP 3 INSECTICIDE

Force® Evo

Insecticide

For Use to Control Insects on Field Corn, Popcorn, Seedcorn, and Sweet Corn

Active Ingredient:	
Tefluthrin*:	24.2%
Other Ingredients:	75.8%
Total:	100.0%

* CAS No. 79538-32-20
Force® Evo is formulated as an emusifiable concentrate (EC) and contains 2.1 pounds tefluthrin per gallon.

KEEP OUT OF REACH OF CHILDREN.



**DANGER / POISON
PELIGRO / VENENO**

ACCEPTED
01/03/2017

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 100-1610

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100- XXXX
EPA Est. XXXX

SCP

2.5 gallons
Net Contents

TABLE OF CONTENTS

1.0 FIRST AID

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

2.2 Personal Protective Equipment (PPE)

2.3 Environmental Hazards

3.0 PRODUCT INFORMATION

3.1 Resistance Management

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

4.2 Application Equipment

4.3 Application Volume and Spray Coverage

4.4 Mixing Directions

5.0 ROTATIONAL CROP RESTRICTIONS

6.0 RESTRICTIONS AND PRECAUTIONS

6.1 Use Restrictions

6.2 Use Precautions

7.0 CROP USE DIRECTIONS

7.1 Corn

8.0 STORAGE AND DISPOSAL

9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

10.0 APPENDIX

10.1 Force Evo Use Summary Table

1.0 FIRST AID

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage. May cause aspiration pneumonia hazard. Contains petroleum distillate.	
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals



DANGER / POISON
PELIGRO / VENENO

Contains Petroleum Distillate. Fatal if absorbed through the skin. Fatal if inhaled. Fatal if swallowed. Corrosive. Causes skin burns. Causes irreversible eye damage. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear coveralls worn over long-sleeved shirt and long pants, socks, chemical-resistant footwear, and chemical-resistant gloves (such as Barrier Laminate \geq 14 mils, Butyl Rubber \geq 14 mils, Nitrile Rubber \geq 14 mils). Remove and wash contaminated clothing before reuse. Do not breathe spray mist. Wear protective eyewear (goggles, face shield, or safety glasses).

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate \geq 14 mils, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or polyvinyl chloride [PVC] \geq 14 mils or Viton® \geq 14 mils
- Wear protective eyewear (goggles, face shield, or safety glasses)
- When mixing and loading wear a chemical resistant apron.
- Chemical-resistant footwear

Important: This product is only to be used in a closed system, handlers must be provided with all PPE specified above and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

2.2.1 USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.2.2 ENGINEERING CONTROL STATEMENTS

Handlers must use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

- Mixers and loaders must use a mechanical transfer system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4).
- The system must be capable of removing the pesticide from the shipping container and transferring it into mixing tanks and/or application equipment.
- At any disconnect point, the system must be equipped with a dry disconnect or dry couple shut-off device that is warranted by the manufacturer to minimize drippage to not more than 2 ml per disconnect point.
- Persons using a closed system that operates under pressure shall wear protective eyewear.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.3 Environmental Hazards

This pesticide is very highly toxic to freshwater and estuarine fish and invertebrates. Use with care when applying in areas adjacent to any body of water. Runoff may be hazardous to aquatic organisms in neighboring areas.

For terrestrial uses: Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Force Evo must be used only in accordance with directions on this label or in separately published Syngenta supplemental labeling for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

Exception: If the product is drenched, soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. No restricted-entry interval (REI) is required following soil injection, soil incorporated or a soil drench application.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate \geq 14 mils, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or polyvinyl chloride [PVC] \geq 14 mils or Viton \geq 14 mils.
- Wear protective eyewear (goggles, face shield, or safety glasses)
- When mixing and loading wear a chemical resistant apron.
- Chemical resistant footwear

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

3.0 PRODUCT INFORMATION

Force Evo is an at-planting, soil-applied insecticide which provides activity against seed-, plant-, and root-feeding insects and helps protect developing corn root systems.

3.0.1 PEST SUPPRESSION

Pest suppression can mean either inconsistent control (good to poor), or consistent control at a level below that generally considered acceptable for commercial control.

3.1 Resistance Management

GROUP 3 INSECTICIDE

Some insects are known to develop resistance to products after repeated use. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label. Consult your local or state agricultural authorities for details.

Force Evo contains a Group 3 insecticide (tefluthrin). Insect biotypes with acquired or inherent resistance to Group 3 insecticides may eventually dominate the insect population if Group 3 insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by Force Evo or other Group 3 insecticides.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

3.1.1 MAINTAINING SUSCEPTIBILITY TO THESE CLASSES OF CHEMISTRY

- Avoid using Group 3 insecticides exclusively for season-long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, apply Force Evo or other Group 3 insecticides using a “treatment window” approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications

(soil, foliar, unless otherwise stated) of the Group 3 insecticides. Do not exceed the maximum Force Evo allowed per year.

- Following a treatment window of Group 3 insecticides, rotate to a treatment window of effective products with a different mode of action before making additional applications of Group 3 insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest's ability to develop resistance to these classes of chemistry.
- If resistance is suspected, do not reapply Force Evo or other Group 3 insecticides.

3.1.2 OTHER INSECT RESISTANCE MANAGEMENT (IRM) PRACTICES

- Incorporate IPM techniques into your insect control program.
- Monitor treated insect populations for loss of field efficacy.
- Use tank-mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied.

3.1.3 OTHER SOURCES FOR INFORMATION ON INSECT RESISTANCE MANAGEMENT

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org/>.

4.0 APPLICATION DIRECTIONS

4.1 Methods of Application

Soil applications of Force Evo are permitted at planting by liquid in-furrow treatment, which ensures a zone of root protection which is approximately 7 inches wide and 5 inches deep, and by liquid T-band treatment as specified in **Section 7.0**.

4.2 Application Equipment

- Spray equipment configuration should be arranged to provide accurate, uniform and thorough delivery of product and minimize potential for spray drift.
- To ensure accuracy, calibrate sprayer before each use.
- For information on spray equipment and calibration, consult spray equipment manufacturers and/or state recommendations.
- All ground application equipment must be properly maintained and calibrated using appropriate carriers.

4.3 Application Volume and Spray Coverage

See **Section 7.0** for application volume information.

4.4 Mixing Directions

This product has been extensively tested by Syngenta and demonstrated to be compatible with liquid in-row starter and pop-up fertilizers and fungicides, which may be applied at-planting simultaneously with this product. Because Force Evo is supplied and used in a closed handling and application system, end users must **not** conduct any additional compatibility tests using a typical jar test. Specific questions about compatibility should be directed to Syngenta personnel.

5.0 ROTATIONAL CROP RESTRICTIONS

The rotational (plant-back) restriction for Force Evo is 30 days for all crops.

6.0 RESTRICTIONS AND PRECAUTIONS

6.1 Use Restrictions

- **DO NOT** apply this product except on the sites and for the pests at the indicated rates and limitations specified in **Section 7.0** of this label.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product within 20 yards of aquatic bodies (lakes, reservoirs, rivers, permanent streams, marshes, estuaries, natural ponds, or commercial fish farm ponds).
 - At least 10 feet of the 20-yard setback area must be left in grass or other natural vegetation.
- **DO NOT** apply this product as a T-band treatment unless the product can be incorporated into the top one inch of soil using tines, chains or other suitable equipment.

6.2 Use Precautions

- When used on highly erodible land (HEL), best management practices for minimizing runoff must be employed.
- When applied to highly erodible land that drains to water bodies, consultation with the Natural Resources Conservation Service (NRCS) and/or other agricultural agencies will ensure that effective efforts are made to minimize soil erosion as well as pesticide runoff.
- When the highly erodible land is adjacent to aquatic bodies, the entire 20-yard buffer/setback must be left to grass or other natural vegetation.
- For NO-TILL corn acreage that drains to adjacent or neighboring aquatic sites, apply as an in-furrow treatment only. See in-furrow directions in **Section 7.0**.

7.0 CROP USE DIRECTIONS

7.1 Corn

Crops (Including all cultivars, varieties, and/or hybrids of these)						
Field Corn	Popcorn		Seed Corn		Sweet Corn	
Target Pest	Rate (fl oz/1000 ft of row)	Application Timing		Use Directions		
Cutworms Lesser Cornstalk Borer Mexican Corn Rootworm Northern Corn Rootworm Seedcorn Beetle Seedcorn Maggot Southern Corn Rootworm Western Corn Rootworm White Grub Wireworm Suppression Only: Billbug Chinch Bug	0.46 – 0.57	Apply at planting.		<p>In-Furrow Applications: Apply into the seed furrow through spray nozzles or micro-tubes, behind the planter furrow openers and in front of the press wheel.</p> <p>Banded Applications: Apply as a 7 inch T-band across the open seed furrow between the furrow openers and the press wheels.</p> <p>Follow the application by an incorporation of 1 inch into the soil either by using tines, press wheels, closing wheels, or other suitable equipment which is either attached to the planter or part of the planter system.</p> <p>Apply in a minimum of 2 gal/A.</p> <p>For best control of Wireworm and/or White Grub, apply the liquid as an in-furrow treatment. For heavy infestations of these pests, use 0.57 fl oz/1000 feet of row.</p> <p>For suppression of Billbug and Chinch Bug, apply as a T-banded application.</p>		
Fluid Ounces of Force Evo Required per Acre for Typical Row Spacings						
Row Spacing	38"	36"	30"	22"	20"	15"
Linear Ft per Acre	13,758	14,520	17,424	23,760	26,136	34,848
Fluid Ounces/1000 ft of row						
0.46 fl. oz	6.5 fl oz	7.0 fl oz	8.0 fl oz	11.0 fl oz	12.0 fl oz	16.0 fl oz
0.57 fl. oz	8.0 fl oz	8.5 fl oz	10.0 fl oz	13.5 fl oz	15.0 fl oz	20.0 fl oz

Resistance Management:

- Refer to **Section 3.1**.

USE RESTRICTIONS

- 1) Refer to **Section 6.1** for additional product use restrictions.
- 2) **Maximum Single Application Rate:** 20.0 fl oz/A (0.327 lb ai/A of tefluthrin-containing products or 0.57 oz/1000 row feet of Force Evo regardless of row spacing)
- 3) **Maximum Annual Rate:** 20.0 fl oz/A/year (0.327 lb ai/A/year of tefluthrin-containing products)
- 4) Make only one application of Force Evo per year.

8.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office.

Container Handling [Bag in a Box]

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

10.0 APPENDIX

10.1 Force Evo Use Summary Table

IMPORTANT: The table below is a summary of the Crop Use Directions for Force Evo. However, it is important for the user to read and follow the complete instructions contained within this label.

Crop or Crop Group or Subgroup, with examples	Maximum Rate per Application (oz/A)	Maximum Rate per Application (lb ai/A)	Minimum Application Interval (days)	Pre-Harvest Interval - PHI (days)	Maximum Rate per Year (oz/A)	Maximum Rate per Year (lb ai/A)
Corn field, pop, seed and sweet	20.0	0.327	NA	NA	20.0	0.327

Force® Evo the ALLIANCE FRAME
the Syngenta Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Viton® is a registered trademark of E.I. DuPont de Nemours and Company

©20XX Syngenta

For non-emergency (e.g. current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP

Force Evo 100-RARN NEW PRODUCT MAY2016 –VER E – mar 12/26/2016
000100-0RARN.20160511E.FORCE_EVO.NEW.MAY2016.PDF

Force Evo 100-RARN NEW PRODUCT MAY2016 –VER E – HI – mar 12/26/2016
000100-0RARN.20160511E.FORCE_EVO.NEW.MAY2016-HI.PDF

Force Evo 100-RARN NEW PRODUCT MAY2016 –VER D – HI – mar 12/20/2016
000100-0RARN.20160511D.FORCE_EVO.NEW.MAY2016-HI.PDF

Force Evo 100-RARN NEW PRODUCT MAY2016 –VER C – HI – mar 12/19/2016
000100-0RARN.20160511C.FORCE_EVO.NEW.MAY2016-HI.PDF

Force Evo 100-XXXX NEW PRODUCT MAY2016 –VER B – HI – mar 12/07/2016
000100-0XXXX.20160511B.FORCE_EVO.NEW.MAY2016-HI.PDF

Force Evo 100-XXXX NEW PRODUCT MAY2016 – mar 5/11/2016
000100-0XXXX.20160511.FORCE_EVO.NEW.MAY2016.PDF